

Gröbner-Shirshov bases for Rota-Baxter algebras*

L. A. Bokut[†]

School of Mathematical Sciences, South China Normal University

Guangzhou 510631, P. R. China

Sobolev Institute of Mathematics, Russian Academy of Sciences

Siberian Branch, Novosibirsk 630090, Russia

bokut@math.nsc.ru

Yuqun Chen[‡] and Xueming Deng

School of Mathematical Sciences, South China Normal University

Guangzhou 510631, P. R. China

yqchen@scnu.edu.cn

dengxm860416@yhao.cn

Abstract: In this paper, we establish the Composition-Diamond lemma for associative nonunitary Rota-Baxter algebras with weight λ . As applications, we obtain a linear basis of a free commutative Rota-Baxter algebra without unity and show that every countably generated Rota-Baxter algebra with weight 0 can be embedded into a two-generated Rota-Baxter algebra.

Key words: Rota-Baxter algebra, Gröbner-Shirshov basis.

AMS 2000 Subject Classification: 16S15, 13P10, 16W99, 17A50

1 Introduction

Gröbner bases and Gröbner-Shirshov bases theories were invented independently by A.I. Shirshov for non-associative algebras and commutative (anti-commutative) non-associative algebras [57], for Lie algebras (explicitly) and associative algebras (implicitly) [58], for infinite series algebras (both formal and convergent) by H. Hironaka [42] and for polynomial

*Supported by the NNSF of China (No.10771077) and the NSF of Guangdong province (No.06025062).

[†]Supported by RFBR 01-09-00157, LSS-344.2008.1 and SB RAS Integration grant No. 2009.97 (Russia).

[‡]Corresponding author.

algebras by B. Buchberger (first publication in [18]). Gröbner bases and Gröbner-Shirshov bases theories have been proved to be very useful in different branches of mathematics, including commutative algebra and combinatorial algebra, see, for example, the books [1, 6, 17, 19, 20, 25, 34], the papers [7, 8, 9], and the surveys [10, 14, 15, 16].

The Shirshov's Composition-Diamond lemma–Buchberger's theorem is the corner stone of the theories. This proposition says that in appropriate free algebra $A_k(X)$ over a field k with a free generating set X and a fixed monomial ordering, the following conditions on a subset S of $A_k(X)$ are equivalent: (i) Any composition (s -polynomial) of polynomials from S is trivial; (ii) If $f \in Id(S)$, then the maximal monomial $\bar{f}$ contains some maximal monomial $\bar{s}$, where $s \in S$; (iii) The set $Irr(S)$ of all (non-associative in general) words in X , which do not contain any maximal word $\bar{s}$, $s \in S$, is a linear k -basis of the algebra $A(X|S) = A(X)/Id(S)$ with generators X and defining relations S (for Lie algebra case, $Irr(S)$ is a set of Lyndon–Shirshov Lie words whose associative support do not contain maximal associative words of polynomials from S).

Up to now, different versions of Composition-Diamond lemma are known for the following classes of algebras apart those mentioned above: (color) Lie super-algebras ([48, 49]) [50], Lie p -algebras [49], associative conformal algebras [13], modules [43], dialgebras [11], associative algebras with multiple operators [12]. In this paper, we establish the Composition-Diamond lemma for associative nonunitary Rota-Baxter algebras with weight λ . As applications, we obtain a linear basis of a free commutative Rota-Baxter algebra without unity and show that every countably generated Rota-Baxter algebra with weight 0 can be embedded into a two-generated Rota-Baxter algebra. In [13], a $\frac{1}{2}$ -PBW Theorem for associative conformal algebras was proved by L.A. Bokut, Y. Fong and W.-F. Ke. Here we prove the similar results for dendriform dialgebra and trialgebra as another two applications of the Composition-Diamond lemma for associative nonunitary Rota-Baxter algebras with weight λ .

Rota-Baxter algebras were invented by G. Baxter [5] and studied by G.-C. Rota [54, 55, 56], F.V. Atkinson [4] and P. Cartier [21]. Since then, it has been related to many topics in mathematics and mathematical physics, see, for example, [22, 23, 24, 26, 27, 28, 29, 32, 33, 35, 36, 40, 41, 44, 51, 52, 53].

Not so many examples of Rota-Baxter algebras are known. They are mostly free associative (commutative and with unity) Rota-Baxter algebras [3, 30, 31] ([37, 38, 41]) and low-dimensional Rota-Baxter algebras [2, 47]. This paper provides a systematical method to deal with Rota-Baxter algebras defined by generators and defining relations.

In this paper, k is a field of characteristic zero and Rota-Baxter algebra always means Rota-Baxter algebra without unity.

We thank Ms Qiuhui Mo for some useful discussions.

2 Free Rota-Baxter algebra

Let A be an associative algebra over k and $\lambda \in k$. Let a k -linear operator $P : A \rightarrow A$ satisfy

$$P(x)P(y) = P(P(x)y) + P(xP(y)) + \lambda P(xy), \forall x, y \in A.$$

Then A is called a Rota-Baxter algebra with weight λ .

A subset I of a Rota-Baxter algebra A is called a Rota-Baxter ideal of A if I is an ideal of A such that $P(I) \subseteq I$.

Let A, B be two Rota-Baxter algebras and $f : A \rightarrow B$ be a mapping. Then f is called a homomorphism if f is a k -algebra homomorphism such that for any $a \in A$, $f(P(a)) = P(f(a))$.

Recall that a free Rota-Baxter algebra with weight λ on a set X is a Rota-Baxter algebra A generated by X with a natural mapping $i : X \rightarrow A$ such that, for any Rota-Baxter algebra R with weight λ and any map $f : X \rightarrow R$, there exists a unique homomorphism $\tilde{f} : A \rightarrow R$ such that $\tilde{f} \cdot i = f$.

The free Rota-Baxter algebra generated by a nonempty set X is given by K. Ebrahimi-Fard and L. Guo [30].

Let X be a nonempty set, $S(X)$ the free semigroup generated by X without identity and P a symbol of a unary operation. For any two nonempty sets Y and Z , denote by

$$\Lambda_P(Y, Z) = (\cup_{r \geq 0} (YP(Z))^r Y) \cup (\cup_{r \geq 1} (YP(Z))^r) \cup (\cup_{r \geq 0} (P(Z)Y)^r P(Z)) \cup (\cup_{r \geq 1} (P(Z)Y)^r),$$

where for a set T , T^0 means the empty set.

Remark: In $\Lambda_P(Y, Z)$, there are no words with a subword $P(z_1)P(z_2)$ where $z_1, z_2 \in Z$.

Define

$$\begin{aligned} \Phi_0 &= S(X) \\ &\vdots \\ \Phi_n &= \Lambda_P(\Phi_0, \Phi_{n-1}) \\ &\vdots \end{aligned}$$

Then

$$\Phi_0 \subset \cdots \subset \Phi_n \subset \cdots$$

Let

$$\Phi(X) = \cup_{n \geq 0} \Phi_n.$$

Clearly, $P(\Phi(X)) \subset \Phi(X)$. If $u \in X \cup P(\Phi(X))$, then u is called prime. For any $u \in \Phi(X)$, u has a unique form $u = u_1 u_2 \cdots u_n$ where u_i is prime, $i = 1, 2, \dots, n$, and u_i, u_{i+1} can not both have forms as $p(u'_i)$ and $p(u'_{i+1})$. If this is the case, then we define the breath of u to be n , denoted by $bre(u) = n$.

For any $u \in \Phi(X)$ and for a set $T \subseteq X \cup \{P\}$, denote by $\deg_T(u)$ the number of occurrences of $t \in T$ in u . Let

$$\text{Deg}(u) = (\deg_{\{P\} \cup X}(u), \deg_{\{P\}}(u)).$$

We order $\text{Deg}(u)$ lexicographically.

Let $k\Phi(X)$ be a free k -module with k -basis $\Phi(X)$ and $\lambda \in k$ a fixed element. Extend linearly $P: k\Phi(X) \rightarrow k\Phi(X)$, $u \mapsto P(u)$ where $u \in \Phi(X)$.

Now we define the multiplication in $k\Phi(X)$.

Firstly, for $u, v \in X \cup P(\Phi(X))$, define

$$u \cdot v = \begin{cases} P(P(u') \cdot v') + P(u' \cdot P(v')) + \lambda P(u' \cdot v'), & \text{if } u = P(u'), v = P(v'); \\ uv, & \text{otherwise.} \end{cases}$$

Secondly, for any $u = u_1 u_2 \cdots u_s, v = v_1 v_2 \cdots v_l \in \Phi(X)$ where u_i, v_j are prime, $i = 1, 2, \dots, s, j = 1, 2, \dots, l$, define

$$u \cdot v = u_1 u_2 \cdots u_{s-1} (u_s \cdot v_1) v_2 \cdots v_l.$$

Equipping with the above concepts, $k\Phi(X)$ is the free Rota-Baxter algebra with weight λ generated by X , see [30].

We denote by $RB(X)$ the free Rota-Baxter algebra with weight λ generated by X .

3 Composition-Diamond lemma for Rota-Baxter algebras

In this section, we establish the Composition-Diamond lemma for Rota-Baxter algebras with weight λ .

Let $\mathcal{N}$ be the set of non-negative integers and $\mathcal{N}^+$ the set of positive integers.

Let the notations be as before. We have to order $\Phi(X)$. Let X be a well ordered set. Let us define an ordering $>$ on $\Phi(X)$ by induction on the Deg -function.

For any $u, v \in \Phi(X)$, if $\text{Deg}(u) > \text{Deg}(v)$, then $u > v$.

If $\text{Deg}(u) = \text{Deg}(v) = (n, m)$, then we define $u > v$ by induction on (n, m) .

If $(n, m) = (1, 0)$, then $u, v \in X$ and we use the ordering on X . Suppose that for (n, m) the ordering is defined where $(n, m) \geq (1, 0)$. Let $(n, m) < (n', m') = \text{Deg}(u) = \text{Deg}(v)$. If $u, v \in P(\Phi(X))$, say $u = P(u')$ and $v = P(v')$, then $u > v$ if and only if $u' > v'$ by induction. Otherwise $u = u_1 u_2 \cdots u_l$ and $v = v_1 v_2 \cdots v_s$ where $l > 1$ or $s > 1$, then $u > v$ if and only if $(u_1, u_2, \dots, u_l) > (v_1, v_2, \dots, v_s)$ lexicographically by induction.

It is clear that $>$ is a well ordering on $\Phi(X)$. Throughout this paper, we will use this ordering.

Now, for any $0 \neq f \in RB(X)$, f has the leading term $\bar{f}$ and $f = \alpha_1 \bar{f} + \sum_{i=2}^n \alpha_i u_i$ where $\bar{f}, u_i \in \Phi(X)$, $\bar{f} > u_i$, $0 \neq \alpha_1, \alpha_i \in k$. Denote by $lc(f)$ the coefficient of the leading term $\bar{f}$. If $lc(f) = 1$, we call f monic.

For any $u \in \Phi(X)$ and any non-negative integer n , we denote by

$$P^n(u) = \underbrace{P(P(\cdots P(u) \cdots))}_n \text{ and } P^0(u) = u.$$

Clearly, if $u \in \Phi(X)$, then there exists a unique $n \in \mathcal{N}$ such that $u = P^n(u')$ where $u' \in \Phi(X) \setminus P(\Phi(X))$.

The proof of the following lemma is straightforward. We omit details.

Lemma 3.1 *For any $u, v \in \Phi(X)$, $n, m \geq 1$, we have*

$$\begin{aligned} & P^n(u) \cdot P^m(v) \\ &= \sum_{s=1}^n \alpha_{(n,m,s)} P^{n+m-s}(P^s(u) \cdot v) + \sum_{l=1}^m \beta_{(n,m,l)} P^{n+m-l}(u \cdot P^l(v)) + \lambda \varepsilon(n, m, u, v) \end{aligned}$$

where $\alpha_{(n,m,s)}, \beta_{(n,m,l)} \in \mathcal{N}^+$, $\varepsilon(n, m, u, v) \in P(RB(X))$, $\deg_{\{P\}}(\overline{\varepsilon(n, m, u, v)}) = \deg_{\{P\}}(P^n(u)) + \deg_{\{P\}}(P^m(v)) - 1$ and $\deg_X(\overline{\varepsilon(n, m, u, v)}) = \deg_X(u) + \deg_X(v)$. Moreover, if we define $\alpha_{(n,0,s)} = 0, \beta_{(0,m,l)} = 0$, then the coefficients satisfy the recursive relations:

- 1) $\alpha_{(n,1,s)} = 1$ and $\beta_{(1,m,l)} = 1$ where $1 \leq s \leq n-1, 1 \leq l \leq m-1$.
- 2) $\alpha_{(n,m,s)} = \alpha_{(n-1,m,s)} + \alpha_{(n,m-1,s)}$ and $\beta_{(n,m,l)} = \beta_{(n-1,m,l)} + \beta_{(n,m-1,l)}$ where $1 \leq s \leq n-1, 1 \leq l \leq m-1$.
- 3) $\alpha_{(n,m,n)} = \beta_{(n,m,m)} = 1$.

Lemma 3.2 *For any $u, v \in \Phi(X)$, if $u, v \notin P(\Phi(X))$, then $\overline{P(u) \cdot v} > \overline{u \cdot P(v)}$. Moreover, for any $m \geq 1, n \geq 1$, $\overline{P^n(u) \cdot P^m(v)} = \overline{P^{n+m-1}(P(u) \cdot v)}$.*

Proof: There are four cases to consider.

Case 1: $bre(u) = bre(v) = 1$, i.e., $u = x, v = y \in X$.

Case 2: $bre(u) = 1$ and $bre(v) = l > 1$, i.e., $u = x \in X, v = v_1 v_2 \cdots v_l$.

Case 3: $bre(u) = s > 1$ and $bre(v) = 1$, i.e., $u = u_1 \cdots u_{s-1} u_s, v = y \in X$.

Since Case 1, Case 2 and Case 3 are simple to prove, we just prove the following.

Case 4: $bre(u) = s > 1$ and $bre(v) = l > 1$, say $u = u_1 \cdots u_{s-1} u_s, v = v_1 v_2 \cdots v_l$. Then we have the following two subcases to consider.

- (i) If $v_1 = y \in X$, then $P(u) \cdot v = P(u_1 \cdots u_{s-1} u_s) y v_2 \cdots v_l, u \cdot P(v) = u_1 \cdots u_{s-1} u_s \cdot P(y v_2 \cdots v_l)$. Since $P(u_1 \cdots u_{s-1} u_s) > u_1$,

$$\overline{P(u) \cdot v} = \overline{P(u_1 \cdots u_{s-1} u_s) y v_2 \cdots v_l} > \overline{u_1 \cdots u_{s-1} u_s \cdot P(y v_2 \cdots v_l)} = \overline{u \cdot P(v)}.$$

- (ii) If $v_1 = P(v'_1)$, then $P(u) \cdot v = (P(u) \cdot P(v'_1))v_2 \cdots v_l$ and $u \cdot P(v) = u_1 \cdots u_{s-1}(u_s \cdot P(v))$.
 Since $\overline{P(u) \cdot P(v'_1)} \in P(\Phi(X))$ and $\overline{P(u) \cdot P(v'_1)} > u_1$,

$$\overline{P(u) \cdot v} = \overline{P(u) \cdot P(v'_1)v_2 \cdots v_l} > u_1 \cdots u_{s-1} \overline{u_s \cdot P(P(v'_1)v_2 \cdots v_l)} = \overline{u \cdot P(v)}. \quad \blacksquare$$

Lemma 3.3 For any $u, v \in \Phi(X)$, if $u > v$, then $\overline{u \cdot w} > \overline{v \cdot w}$ and $\overline{w \cdot u} > \overline{w \cdot v}$ for any $w \in \Phi(X)$.

Proof: Firstly, we show $\overline{u \cdot w} > \overline{v \cdot w}$. We may assume $u = u_1 \cdots u_s, v = v_1 \cdots v_l, s, l \geq 1$ and $u_1 > v_1$ where u_i, v_j are prime, $i = 1, 2, \dots, s, j = 1, 2, \dots, l$, and $\text{Deg}(u) = \text{Deg}(v)$. We may also assume that $w = P(w') \in P(\Phi(X))$.

It is noted that if $s > 1$ and $l = 1$, then, since $\text{Deg}(u) = \text{Deg}(v)$, we will have $v > u$, a contradiction. So we just need to consider the following cases.

Case 1: $s > 1, l > 1$ and Case 2: $s = 1, l > 1$. These two cases are clear.

Case 3: $s = l = 1$. If $u_1 \in X$, then $v_1 \in X$ and the result holds clearly. If $u_1 \in P(\Phi(X))$, then $v_1 \in P(\Phi(X))$. We may assume that $u_1 = P^n(u'_1), v_1 = P^{n'}(v'_1)$ with $u'_1, v'_1 \notin P(\Phi(X))$ and $w = P^m(w''), w'' \notin P(\Phi(X))$. Then we need to consider two subcases.

- (i) If $n > n'$, then $\overline{u \cdot w} = P^{n+m-1}(\overline{P(u'_1) \cdot w''}) > P^{n'+m-1}(\overline{P(v'_1) \cdot w''}) = \overline{v \cdot w}$ by Lemma 3.2.

- (ii) If $n = n'$, then $u'_1 > v'_1$. By Lemma 3.2 we have

$$\overline{u \cdot w} = P^{n+m-1}(\overline{P(u'_1) \cdot w''}), \quad \overline{v \cdot w} = P^{n+m-1}(\overline{P(v'_1) \cdot w''}).$$

By induction on $\text{deg}_{\{P\} \cup X}(w)$, $\overline{P(u'_1) \cdot w''} > \overline{P(v'_1) \cdot w''}$ and so $\overline{u \cdot w} > \overline{v \cdot w}$.

This shows that $\overline{u \cdot w} > \overline{v \cdot w}$.

Secondly, we show that $\overline{w \cdot u} > \overline{w \cdot v}$.

We may assume that $w = P(w') \in P(\Phi(X))$, $\text{Deg}(u) = \text{Deg}(v)$, $u = u_1 u_2 \cdots u_s > v = v_1 v_2 \cdots v_l$ where u_i, v_j are prime, $i = 1, 2, \dots, s, j = 1, 2, \dots, l$. Then we have the following two cases.

Case 1: $u_1 > v_1$. We only need to consider the case of $u_1 = P^n(u'_1) \in P(\Phi(X))$, $v_1 = P^{n'}(v'_1) \in P(\Phi(X))$ where $n, n' \geq 1, u'_1, v'_1 \notin P(\Phi(X))$. Let $w = P^m(w'') \in P(\Phi(X))$ where $m \geq 1$ and $w'' \notin P(\Phi(X))$. Then we have the following subcases to consider.

- (i) If $n > n'$, then $\overline{w \cdot u} = P^{n+m-1}(\overline{P(w'') \cdot u'_1})u_2 \cdots u_s > P^{n'+m-1}(\overline{P(w'') \cdot v'_1})v_2 \cdots v_l = \overline{w \cdot v}$ by Lemma 3.2.
- (ii) If $n = n'$, then $u'_1 > v'_1$. By induction on $\text{deg}_{\{P\} \cup X}(u) + \text{deg}_{\{P\} \cup X}(v)$, we have $\overline{P(w'') \cdot u'_1} > \overline{P(w'') \cdot v'_1}$. Therefore, by Lemma 3.2

$$\overline{w \cdot u} = P^{n+m-1}(\overline{P(w'') \cdot u'_1})u_2 \cdots u_s > P^{n+m-1}(\overline{P(w'') \cdot v'_1})v_2 \cdots v_l = \overline{w \cdot v}.$$

Case 2: $u_1 = v_1$. This case is clear. ■

For any $u, v \in \Phi(X)$, clearly,

$$u > v \implies P(u) > P(v).$$

Let $\star$ be a symbol and $\star \notin X$. By a $\star$ -Rota-Baxter word we mean any expression in $\Phi(X \cup \{\star\})$ with only one occurrence of $\star$. The set of all $\star$ -Rota-Baxter words on X is denoted by $\Phi^\star(X)$.

Let u be a $\star$ -Rota-Baxter word and $s \in RB(X)$. Then we call

$$u|_s = u|_{\star \mapsto s}$$

an s -Rota-Baxter word. For short, we call $u|_s$ an s -word.

In other words, an s -word $u|_s$ means that we have replaced the $\star$ of u by s .

For example, if $u = P(x_1)x_2P^2(\star)x_4P(x_5)$, then $u|_s = P(x_1)x_2P^2(s)x_4P(x_5)$ is an s -word.

Similarly, we can define $(\star_1, \star_2)$ -Rota-Baxter words as expressions in $\Phi(X \cup \{\star_1, \star_2\})$ with only one occurrence of $\star_1$ and only one occurrence of $\star_2$. Let us denote by $\Phi^{\star_1, \star_2}(X)$ the set of all $(\star_1, \star_2)$ -Rota-Baxter words. Let $u \in \Phi^{\star_1, \star_2}(X)$. Then we call

$$u|_{s_1, s_2} = u|_{\star_1 \mapsto s_1, \star_2 \mapsto s_2}$$

an s_1 - s_2 -Rota-Baxter word. For short, we call $u|_{s_1, s_2}$ an s_1 - s_2 -word.

If $\overline{u|_s} = u|_{\overline{s}}$, then we call $u|_s$ a normal s -word.

By Lemma 3.3, we have the following lemma which shows that the ordering $>$ on $\Phi(X)$ is monomial.

Lemma 3.4 *For any $u, v \in \Phi(X)$, $w \in \Phi^\star(X)$,*

$$u > v \implies \overline{w|_u} > \overline{w|_v}$$

where $w|_u = w|_{\star \mapsto u}$ and $w|_v = w|_{\star \mapsto v}$. ■

Remark: If $u|_s$ is a normal s -word, then $P^l(u|_s)$ is also a normal s -word where $l \in \mathcal{N}$.

In order to describe the ideal $Id(S)$ of $RB(X)$ generated by S , we introduce the concept of P - s -words.

For any nonempty set X , define

$$\begin{aligned} \Psi_0 &= S(X) \\ \Psi_1 &= S(X \cup P(S(X))) \\ &\vdots \\ \Psi_n &= S(X \cup P(\Psi_{n-1})) \\ &\vdots \end{aligned}$$

Then we have

$$\Psi_0 \subset \Psi_1 \subset \cdots \subset \Psi_n \subset \cdots$$

Let

$$\Psi(X) = \cup_{n \geq 0} \Psi_n.$$

Clearly, $\Phi(X) \subset \Psi(X)$.

Now, we can define $\Psi^*(X)$ as the definition of $\Phi^*(X)$.

Let $S \subseteq RB(X)$. Then it is clear that $Id(S) = span_k\{w|_s \mid w \in \Psi^*(X), s \in S\}$ where $span_k\{w|_s \mid w \in \Psi^*(X), s \in S\}$ is the k -subspace of $RB(X)$ generated by $\{w|_s \mid w \in \Psi^*(X), s \in S\}$.

Let $w \in \Psi^*(X)$ and $s \in S$. Then we call $w|_s$ a P - s -word.

The following lemma says that each P - s -word is a linear combination of s -words.

Lemma 3.5 *Let $S \subseteq RB(X)$, $u' \in \Psi^*(X)$ and $s \in S$. Then the P - s -word $u'|_s$ has an expression:*

$$u'|_s = \sum \alpha_i u_i|_s, \quad \text{where each } \alpha_i \in k, \quad u_i|_s \text{ is } s\text{-word and } \overline{u_i|_s} \leq \overline{u'|_s}.$$

In particular, $Id(S) = span_k\{w|_s \mid w \in \Phi^*(X), s \in S\}$.

Proof: For any $u \in \Psi(X)$, in $\Psi(X)$, u has a unique expression

$$u = u_1 \cdots u_n, \quad u_i \in X \cup P(\Psi(X)).$$

Such an n is called the P -length of u .

Let $u'|_s$ be a P - s -word. Denote by $u'' = u'|_{\star \mapsto x} = u_1 \cdots u_n$, where $x \in X$, $u_i \in X \cup P(\Psi(X))$. We prove the result by induction on the P -length n of u'' and on $\overline{u'|_s}$.

We may assume that $u'' \notin \Phi(X)$.

Case 1. $n = 1$. Then $u'|_s = P(u|_s)$. Since $\overline{u|_s} < \overline{u'|_s}$, the result follows by induction on $\overline{u'|_s}$.

Case 2. $n \geq 2$. Then there are two cases to consider.

2.1 There exists $1 \leq i \leq n-1$ such that $u_i = P(u'_i)$ and $u_{i+1} = P(u'_{i+1})$ where $u'_i, u'_{i+1} \in \Psi(X)$. Thus, by the Rota-Baxter identity, in $RB(X)$,

$$u'' = u_1 \cdots u_{i-1} (P(P(u'_i)u'_{i+1}) + P(u'_i P(u'_{i+1})) + \lambda P(u'_i u'_{i+1})) u_{i+2} \cdots u_n,$$

where in the right side, each term has P -length $n-1$ and $\overline{(u_1 \cdots u_{i-1} P(P(u'_i)u'_{i+1})u_{i+2} \cdots u_n)|_s}, \overline{(u_1 \cdots u_{i-1} P(u'_i P(u'_{i+1}))u_{i+2} \cdots u_n)|_s}, \overline{(u_1 \cdots u_{i-1} P(u'_i u'_{i+1})u_{i+2} \cdots u_n)|_s} \leq \overline{u'|_s}$. Now the result follows by induction on n .

2.2 For each j , $1 \leq j \leq n-1$, $\{u_j, u_{j+1}\} \cap X \neq \emptyset$. Let $u'|_s = u_1 \cdots u_{m-1} u_m|_s u_{m+1} \cdots u_n$.

For u_m , there are two cases to consider: $u_m = P(v)$ and $u_m = x$.

For each u_i , $1 \leq i \leq n$, $i \neq m$, by the Rota-Baxter identity, we may assume

$$u_i = \sum \alpha_{iq_i} v_{iq_i}, \quad \text{where each } \alpha_{iq_i} \in k, \quad v_{iq_i} \in \Phi(X), \quad v_{iq_i} \leq \overline{u_i}.$$

Thus,

$$u_1 \cdots u_m \cdots u_n = \sum \alpha_{1q_1} \cdots \widehat{\alpha_{mq_m}} \cdots \alpha_{nq_n} v_{1q_1} \cdots u_m \cdots v_{nq_n}$$

where each $v_{1q_1} \cdots y \cdots v_{nq_n} \in \Phi^*(X)$, $y \in \{P(\star), \star\}$, $v_{1q_1} \cdots \overline{u_m} \cdots v_{nq_n} \leq \overline{u_1 \cdots u_m \cdots u_n}$, and $v_{1q_1} \cdots u_m \cdots v_{nq_n}$ has P -length n .

If $u_m = P(v)$ then the result follows from the induction on $\overline{u'|_s}$ since $\overline{v|_s} < \overline{u'|_s}$.

If $u_m = x$ then the result is clear.

The proof is complete. ■

Let $f, g \in RB(X)$ be monic with $\overline{f} = u_1 u_2 \cdots u_n$ where each u_i is prime. Then, there are four kinds of compositions.

- (i) If $u_n \in P(\Phi(X))$, then we define composition of right multiplication as $f \cdot u$ where $u \in P(\Phi(X))$.
- (ii) If $u_1 \in P(\Phi(X))$, then we define composition of left multiplication as $u \cdot f$ where $u \in P(\Phi(X))$.
- (iii) If there exists a $w = \overline{f}a = b\overline{g}$ where fa is normal f -word and bg is normal g -word, $a, b \in \Phi(X)$ and $\deg_{\{P\} \cup X}(w) < \deg_{\{P\} \cup X}(\overline{f}) + \deg_{\{P\} \cup X}(\overline{g})$, then we define the intersection composition of f and g with respect to w as $(f, g)_w = f \cdot a - b \cdot g$.
- (iv) If there exists a $w = \overline{f} = u|_{\overline{g}}$ where $u \in \Phi^*(X)$, then we define the inclusion composition of f and g with respect to w as $(f, g)_w = f - u|_g$. Note that if this is the case, then $u|_g$ is a normal g -word.

We call w in $(f, g)_w$ the ambiguity with respect to f and g . By Lemma 3.4,

$$\overline{(f, g)_w} < w.$$

Let $S \subset RB(X)$ be a set of monic polynomials. Then the composition $(f, g)_w$ is called trivial modulo (S, w) if

$$(f, g)_w = \sum_i \alpha_i u_i|_{s_i}$$

where each $\alpha_i \in k$, $s_i \in S$, $u_i|_{s_i}$ is normal s_i -word and $u_i|_{\overline{s_i}} < w$. If this is the case, then we write

$$(f, g)_w \equiv 0 \pmod{(S, w)}.$$

The composition of left (right) multiplication is called trivial $\text{mod}(S)$ if

$$u \cdot f = \sum_i \alpha_i u_i|_{s_i} \quad (f \cdot u = \sum_i \alpha_i u_i|_{s_i})$$

where each $\alpha_i \in k$, $s_i \in S$, $u_i|_{s_i}$ is normal s_i -word and $u_i|_{\overline{s_i}} \leq \overline{u \cdot f}$ ($u_i|_{\overline{s_i}} \leq \overline{f \cdot u}$). If this is the case, then we write

$$u \cdot f \equiv 0 \pmod{S} \quad (f \cdot u \equiv 0 \pmod{S}).$$

In general, for any two polynomials p and q , $p \equiv q \pmod{S, w}$ means that $p - q = \sum_i \alpha_i u_i|_{s_i}$ where each $\alpha_i \in k$, $s_i \in S$, $u_i|_{s_i}$ is normal s_i -word and $u_i|_{\overline{s_i}} < w$.

S is called a Gröbner-Shirshov basis in $RB(X)$ if each composition $(f, g)_w$ of $f, g \in S$ is trivial $\text{mod}(S, w)$ and each composition of left multiplication and right multiplication is trivial $\text{mod}(S)$.

Lemma 3.6 *Let $S \subset RB(X)$ be a set of monic polynomials. If each composition of left multiplication and right multiplication of S is trivial $\text{mod}(S)$, then each s -word $u|_s$ has an expression*

$$u|_s = \sum_i \alpha_i u_i|_{s_i}$$

where each $\alpha_i \in k$, $s_i \in S$, $u_i|_{s_i}$ is normal s_i -word and $u_i|_{\overline{s_i}} \leq \overline{u|_s}$.

Proof: We prove the result by induction on $\overline{u|_s}$. If $\overline{u|_s} = \overline{s}$, then there is nothing to prove. Suppose that $\overline{u|_s} > \overline{s}$. We consider the following two cases.

Case 1: $u|_s = w_1 s w_2$ where $w_1, w_2 \in \Phi(X)$ (w_1, w_2 may be empty). We only consider the case $w_1, w_2 \in \Phi(X)$. The other cases are simple. By induction we assume that $w_1 s = \sum_i \alpha_i v_i|_{s_i}$ where each $v_i|_{s_i}$ is a normal s_i -word, $s_i \in S$, $\alpha_i \in k$ and $v_i|_{\overline{s_i}} \leq \overline{w_1 s}$.

Suppose that $w_2 = w_{21} w_{22} \cdots w_{2n}$ and $\overline{s_i} = q_{i1} q_{i2} \cdots q_{it}$ where w_{2j}, q_{ij} are prime.

There are two cases to consider.

(I) If $v_i|_{s_i} = w_{1i} s_i w_{2i}$, then we need to consider the following four subcases.

Subcase 1: w_{1i} and w_{2i} are both empty words. Then

$$v_i|_{s_i} w_2 = s_i w_2.$$

In this case, we also have the following two subcases to consider.

(I-1-1): $q_{it} \in X$. In this case, $v_i|_{s_i} w_2 = s_i w_2$ is clearly a normal s_i -word.

(I-1-2): $q_{it} \in P(\Phi(X))$. If $w_{21} \in X$, then $v_i|_{s_i} w_2 = s_i w_2$ is normal s_i -word clearly.

Otherwise, since each composition of right multiplication is trivial,

$$s_i w_2 = \sum_j \gamma_{ij} v'_{ij}|_{s_{ij}} \quad (*)$$

where each $\gamma_{ij} \in k$, $s_{ij} \in S$, $v'_{ij}|_{s_{ij}}$ is normal s_{ij} -word and $v'_{ij}|_{\overline{s_{ij}}} \leq \overline{s_i w_2} = \overline{v_i|_{s_i} w_2} \leq \overline{w_1 s w_2} = \overline{u|_s}$.

Subcase 2: $w_{1i} \in \Phi(X)$ and w_{2i} is empty word. Then

$$v_i|_{s_i} w_2 = w_{1i} s_i w_2.$$

If $\text{bre}(\overline{s_i}) = 1$, then we have the following two subcases to consider.

(I-2-1): $\overline{s_i} = x \in X$. Then $v_i|_{s_i} w_2$ is a normal s_i -word clearly.

(I-2-2): $\overline{s_i} \in P(\Phi(X))$. Then we need to consider the following two subcases.

(i): If $w_{21} \in X$, then $v_i|_{s_i} w_2$ is a normal s_i -word.

(ii): If $w_{21} \in P(\Phi(X))$, then because each composition of right multiplication is trivial, we have (*). Since $v_i|_{s_i} = w_{1i} s_i$ is normal s_i -word and $\overline{s_i} \in P(\Phi(X))$, we have $w_{1i} = w'_{1i} x$ where $x \in X$. Therefore, each $w_{1i} v'_{ij}|_{s_{ij}}$ is still normal s_{ij} -word. Thus, $v_i|_{s_i} w_2 = \sum_j \beta_{ij} w_{1i} v'_{ij}|_{s_{ij}}$ where each $u_{ij}|_{s_{ij}} = \overline{w_{1i} v'_{ij}|_{s_{ij}}}$ is normal s_{ij} -word and $u_{ij}|\overline{s_{ij}} = w_{1i} v'_{ij}|\overline{s_{ij}} \leq \overline{w_{1i} \overline{s_i} w_2} = \overline{v_i|_{s_i} w_2} \leq \overline{u|_s}$.

If $\text{bre}(\overline{s_i}) = t > 1$, we also have the following two subcases to consider.

(I-2-3): $q_{it} \in X$. In this case, $v_i|_{s_i} w_2 = w_{1i} s_i w_2$ is clearly a normal s_i -word.

(I-2-4): $q_{it} \in P(\Phi(X))$. If $w_{21} \in X$, then $v_i|_{s_i} w_2 = w_{1i} s_i w_2$ is normal s_i -word clearly. Otherwise, since each composition of right multiplication is trivial, we have (*). We need to consider the following two cases.

(i): $q_{i1} = P(q'_{i1})$. Similar to the case (ii) in (I-2-2), we have the result.

(ii): $q_{i1} \in X$. We need to consider the following two subcases.

(a): If $v'_{ij}|\overline{s_{ij}} < \overline{s_i w_2}$, then $\overline{w_{1i} v'_{ij}|_{s_{ij}}} < \overline{v_i|_{s_i} w_2} \leq \overline{u|_s}$. By induction, $w_{1i} v'_{ij}|_{s_{ij}} = \sum_m \alpha_{ijm} u_{ijm}|_{s_{ijm}}$ where each $s_{ijl} \in S$, $u_{ijm}|_{s_{ijm}}$ is normal s_{ijm} -word and $u_{ijm}|\overline{s_{ijm}} \leq \overline{w_{1i} v'_{ij}|_{s_{ij}}} < \overline{u|_s}$.

(b): If $v'_{ij}|\overline{s_{ij}} = \overline{s_i w_2}$, because $\overline{s_i w_2} = q_{i1} \cdots q_{i(t-1)} \overline{q_{it} w_2}$ and $q_{i1} \in X$, we have $w_{1i} v'_{ij}|_{s_{ij}}$ is still normal s_{ij} -word.

Subcase 3: $w_{1i}, w_{2i} \in \Phi(X)$ and Subcase 4: w_{1i} is a empty word and $w_{2i} \in \Phi(X)$. These two subcases are simple to prove.

(II) $v_i|_{s_i} = w_{1i} P(v'_i|_{s_i}) w_{2i}$. Since $v_i|_{s_i}$ is a normal s_i word, we may just consider the case w_{1i} and w_{2i} are both empty word (other cases are simple). Moreover, we may suppose $w_2 = P(w'_2)$. Then we have

$$v_i|_{s_i} w_2 = P(v'_i|_{s_i}) \cdot P(w'_2) = P(P(v'_i|_{s_i}) \cdot w'_2) + P(v'_i|_{s_i} \cdot P(w'_2)) + \lambda P(v'_i|_{s_i} \cdot w'_2).$$

Similar to the proof of Lemma 3.5, we have the result.

This proves the Case 1.

Case 2: $u|_s = w_1 P(u'|_s) w_2$ where $u' \in \Phi^*(X)$, $w_1, w_2 \in \Phi(X)$ (w_1, w_2 may be empty). By induction we have $u'|_s = \sum_i \alpha_i u'_i|_{s_i}$ where each $s_i \in S$, $u'_i|_{s_i}$ is normal s_i -word and $\overline{u'_i|_{s_i}} \leq \overline{u'|_s}$. Thus $u|_s = \sum_i \alpha_i \overline{u'_i|_{s_i}}$ where each $u_i|_{s_i} = w_1 P(u'_i|_{s_i}) w_2$ is normal s_i -word and $u_i|\overline{s_i} = w_1 P(u'_i|\overline{s_i}) w_2 \leq \overline{w_1 P(u'|_s) w_2} = \overline{u|_s}$. ■

Lemma 3.7 *Let S be a Gröbner-Shirshov basis in $RB(X)$, $u_1, u_2 \in \Phi^*(X)$ and $s_1, s_2 \in S$. If $w = u_1|_{\overline{s_1}} = u_2|_{\overline{s_2}}$ where each $u_i|_{s_i}$ is a normal s_i -word, $i = 1, 2$, then*

$$u_1|_{s_1} \equiv u_2|_{s_2} \pmod{(S, w)}.$$

Proof: There are three cases to consider. In fact, all cases are essentially true in the same way.

(i) $\overline{s_1}$ and $\overline{s_2}$ are disjoint. In this case, there exists a $(\star_1, \star_2)$ -Rota-Baxter word Π such that

$$\Pi|_{\overline{s_1}, \overline{s_2}} = u_1|_{\overline{s_1}} = u_2|_{\overline{s_2}}$$

where $u_1|_{s_1} = \Pi|_{s_1, \overline{s_2}}$ is normal s_1 -word and $u_2|_{s_2} = \Pi|_{\overline{s_1}, s_2}$ is normal s_2 -word. Then

$$u_2|_{s_2} - u_1|_{s_1} = \Pi|_{\overline{s_1}, s_2} - \Pi|_{s_1, \overline{s_2}} = -\Pi|_{s_1 - \overline{s_1}, s_2} + \Pi|_{s_1, s_2 - \overline{s_2}}.$$

By Lemma 3.5, we may suppose that

$$-\Pi|_{s_1 - \overline{s_1}, s_2} = \sum_t \alpha_{2t} u_{2t}|_{s_2}, \quad \Pi|_{s_1, s_2 - \overline{s_2}} = \sum_l \alpha_{1l} u_{1l}|_{s_1},$$

where $u_{2t}, u_{1l} \in \Phi^*(X)$.

Since S is a Gröbner-Shirshov basis, by Lemma 3.6, we have

$$u_{2t}|_{s_2} = \sum_n \beta_{2tn} w_{2tn}|_{s_{2tn}}$$

and

$$u_{1l}|_{s_1} = \sum_m \beta_{1lm} w_{1lm}|_{s_{1lm}}$$

where $\beta_{1lm}, \beta_{2tn} \in k$, $s_{2tn}, s_{1lm} \in S$, $w_{2tn}|_{s_{2tn}}$ and $w_{1lm}|_{s_{1lm}}$ are normal s_{2tn} - and s_{1lm} -words, respectively, $w_{2tn}|_{\overline{s_{2tn}}} \leq u_{2t}|_{s_2} = \Pi|_{s_1 - \overline{s_1}, s_2} < \Pi|_{\overline{s_1}, s_2} = w$ and $w_{1lm}|_{\overline{s_{1lm}}} < w$. Therefore,

$$u_2|_{s_2} - u_1|_{s_1} = \sum_{t,n} \alpha_{2t} \beta_{2tn} w_{2tn}|_{s_{2tn}} + \sum_{l,m} \alpha_{1l} \beta_{1lm} w_{1lm}|_{s_{1lm}}.$$

It follows that

$$u_1|_{s_1} \equiv u_2|_{s_2} \pmod{(S, w)}.$$

(ii) $\overline{s_1}$ and $\overline{s_2}$ have nonempty intersection but do not include each other. Without loss of generality, we may assume that $\overline{s_1}a = b\overline{s_2}$ for some $a, b \in \Phi(X)$. This implies that s_1a is normal s_1 -word and bs_2 is normal s_2 -word. Then there exists a $\Pi \in \Phi^*(X)$ such that

$$\Pi|_{\overline{s_1}a} = u_1|_{\overline{s_1}} = u_2|_{\overline{s_2}} = \Pi|_{b\overline{s_2}}$$

where $\Pi|_{s_1a}$ is normal s_1a -word and $\Pi|_{bs_2}$ is normal bs_2 -word. Thus, we have

$$u_2|_{s_2} - u_1|_{s_1} = \Pi|_{bs_2} - \Pi|_{s_1a} = -\Pi|_{s_1a - bs_2}.$$

Since S is a Gröbner-Shirshov basis,

$$s_1 a - b s_2 = \sum_j \alpha_j v_j|_{s_j}$$

where each $\alpha_j \in k$, $s_j \in S$, $v_j|_{\overline{s_j}} < \overline{s_1} a$ and $v_j|_{s_j}$ is normal s_j -word. Let $\Pi|_{v_j|_{s_j}} = \Pi_j|_{s_j}$. Then, by Lemmas 3.5 and 3.6,

$$\Pi_j|_{s_j} = \sum_l \beta_{jl} w_{jl}|_{s_{jl}}$$

where each $\beta_{jl} \in k$, $s_{jl} \in S$, $w_{jl}|_{s_{jl}}$ is also normal s_{jl} -word, $w_{jl}|_{\overline{s_{jl}}} \leq \overline{\Pi_j|_{s_j}} = \overline{\Pi|_{v_j|_{s_j}}} < \Pi|_{\overline{s_1} a} = w$. Therefore

$$u_2|_{s_2} - u_1|_{s_1} = \sum_j \alpha_j \Pi_j|_{s_j} = \sum_{j,l} \alpha_j \beta_{jl} w_{jl}|_{s_{jl}}.$$

It follows that

$$u_1|_{s_1} \equiv u_2|_{s_2} \pmod{(S, w)}.$$

(iii) One of $\overline{s_1}$, $\overline{s_2}$ is contained in the other. For example, let $\overline{s_1} = u|_{\overline{s_2}}$ for some $\star$ -Rota-Baxter word u . Then

$$w = u_2|_{\overline{s_2}} = u_1|_{u|_{\overline{s_2}}}$$

and

$$u_2|_{s_2} - u_1|_{s_1} = u_1|_{u|_{s_2}} - u_1|_{s_1} = -u_1|_{s_1 - u|_{s_2}}.$$

Similar to (ii), we can obtain the result. ■

Lemma 3.8 *Let $S \subseteq RB(X)$ be a set of monic polynomials. $Irr(S) \triangleq \{u \in \Phi(X) | u \neq v|_{\overline{s}}, s \in S, v|_{\overline{s}} \text{ is normal } s\text{-word}\}$. Then for any $f \in RB(X)$, f has an expression*

$$f = \sum_i \alpha_i u_i + \sum_j \beta_j v_j|_{s_j}$$

where $\alpha_i, \beta_j \in k$, $u_i \in Irr(S)$, $\overline{u_i} \leq \overline{f}$, $s_j \in S$, $v_j|_{s_j}$ is normal s_j -word and $v_j|_{\overline{s_j}} \leq \overline{f}$.

Proof: Let $f = \sum_i \alpha_i u_i$, $0 \neq \alpha_i \in k, u_1 > u_2 > \dots$. If $u_1 \in Irr(S)$, then let $f_1 = f - \alpha_1 u_1$. If $u_1 \notin Irr(S)$, i.e., there exists $s_1 \in S$ such that $u_1 = v_1|_{\overline{s_1}}$ where $v_1|_{s_1}$ is normal s_1 -word, then let $f_1 = f - \alpha_1 v_1|_{s_1}$. In both cases $\overline{f_1} < \overline{f}$. Now, the result follows from induction on $\overline{f}$. ■

Theorem 3.9 *(Composition-Diamond lemma for Rota-Baxter algebras) Let $RB(X)$ be a free Rota-Baxter algebra over a field of characteristic 0 and S a set of monic polynomials in $RB(X)$, $>$ the monomial ordering on $\Phi(X)$ defined as before and $Id(S)$ the Rota-Baxter ideal of $RB(X)$ generated by S . Then the following statements are equivalent.*

(I) S is a Gröbner-Shirshov basis in $RB(X)$.

(II) $f \in Id(S) \Rightarrow \bar{f} = u|_{\bar{s}}$ for some $u \in \Phi^*(X)$, $s \in S$.

(III) $Irr(S) = \{u \in \Phi(X) | u \neq v|_{\bar{s}}, s \in S, v|_s \text{ is normal } s\text{-word}\}$ is a k -basis of $RB(X|S)$
 $= RB(X)/Id(S)$.

Proof: (I) $\Rightarrow$ (II) Let $0 \neq f \in Id(S)$. By Lemmas 3.5 and 3.6, we may assume that $f = \sum_{i=1}^n \alpha_i u_i|_{s_i}$ where each $\alpha_i \in k$, $s_i \in S$ and $u_i|_{s_i}$ is normal s_i -word. Let $w_i = u_i|_{\bar{s}_i}$. We arrange these leading words in non-increasing order by

$$w_1 = w_2 = \cdots = w_m > w_{m+1} \geq \cdots \geq w_n.$$

Now we prove the result by induction on m and w_1 .

If $m = 1$, then $\bar{f} = u_1|_{\bar{s}_1}$ and there is nothing to prove. Now we assume that $m \geq 2$. Then $u_1|_{\bar{s}_1} = w_1 = w_2 = u_2|_{\bar{s}_2}$. Since S is a Gröbner-Shirshov basis in $RB(X)$, by Lemma 3.7, we have

$$u_2|_{s_2} - u_1|_{s_1} = \sum_j \beta_j v_j|_{s_j}$$

where each $\beta_j \in k$, $s_j \in S$, $v_j|_{\bar{s}_j} < w_1$ and $v_j|_{s_j}$ is normal s_j -word. Therefore, since

$$\alpha_1 u_1|_{s_1} + \alpha_2 u_2|_{s_2} = (\alpha_1 + \alpha_2) u_1|_{s_1} + \alpha_2 (u_2|_{s_2} - u_1|_{s_1}),$$

we have

$$f = (\alpha_1 + \alpha_2) u_1|_{s_1} + \sum_j \alpha_2 \beta_j v_j|_{s_j} + \sum_{i=3}^n \alpha_i u_i|_{s_i}.$$

If either $m > 2$ or $\alpha_1 + \alpha_2 \neq 0$, then the result follows from induction on m . If $m = 2$ and $\alpha_1 + \alpha_2 = 0$, then the result follows from induction on w_1 .

(II) $\Rightarrow$ (III) For any $f \in RB(X)$, by Lemma 3.8, $f + Id(S)$ can be expressed by the elements of $Irr(S) + Id(S)$. Now Suppose that $\alpha_1 u_1 + \alpha_2 u_2 + \cdots + \alpha_n u_n = 0$ in $RB(X|S)$ with each $u_i \in Irr(S)$, $u_1 > u_2 > \cdots > u_n$ and each $0 \neq \alpha_i \in k$. Then, in $RB(X)$,

$$g = \alpha_1 u_1 + \alpha_2 u_2 + \cdots + \alpha_n u_n \in Id(S).$$

By (II), we have $u_1 = \bar{g} \notin Irr(S)$, a contradiction. Hence, $Irr(S)$ is k -linearly independent. This shows that $Irr(S)$ is a k -basis of $RB(X|S)$.

(III) $\Rightarrow$ (I) For any $(f, g)_w$ where $f, g \in S$, by Lemma 3.8, $(f, g)_w = \sum_i \alpha_i u_i + \sum_j \beta_j v_j|_{s_j}$ where $\alpha_i, \beta_j \in k$, $u_i \in Irr(S)$, $s_j \in S$, $\bar{u}_i \leq \overline{(f, g)_w} < w$, $v_j|_{s_j}$ is normal s_j -word and $v_j|_{\bar{s}_j} \leq \overline{(f, g)_w} < w$. Since $(f, g)_w \in Id(S)$ and by (III), each $\alpha_i = 0$. Thus $(f, g)_w \equiv 0 \pmod{(S, w)}$.

For any composition of left multiplication, $P(u) \cdot f$ where $f \in S$, by Lemma 3.8 and by (III), we have $P(u) \cdot f = \sum_j \beta_j v_j|_{s_j}$ where each $\beta_j \in k$, $s_j \in S$, $v_j|_{s_j}$ is normal s_j -word and $v_j|_{\bar{s}_j} \leq \overline{P(u) \cdot f}$. This shows that each composition of left multiplication in S is trivial $\pmod{(S)}$. Similarly, each composition of right multiplication in S is trivial $\pmod{(S)}$. ■

4 Applications

In [54, 21], G. Rota and P. Cartier constructed free commutative Rota-Baxter algebra with certain restriction. While in [37, 38], L. Guo and W. Keigher constructed free commutative Rota-Baxter algebra with unity by mixable shuffle algebra, also see [41]. In this section, by using the Composition-Diamond lemma for Rota-Baxter algebras (Theorem 3.9), we have the following results: 1. We give a Gröbner-Shirshov basis for the free commutative Rota-Baxter algebra $RB(X|S)$ and as an application, a linear basis of the free commutative Rota-Baxter algebra without unity is obtained. 2. We show that every countably generated Rota-Baxter algebra with weight 0 can be embedded into a two-generated Rota-Baxter algebra. 3. We prove the $\frac{1}{2}$ -PBW Theorems for dendriform dialgebra and dendriform trialgebra.

Theorem 4.1 *Let I be a well ordered set, $X = \{x_i | i \in I\}$ and the ordering on $\Phi(X)$ defined as before. Let*

$$f = x_i x_j - x_j x_i, \quad i > j, \quad i, j \in I \quad (1)$$

$$g = P(u)x_i - x_i P(u), \quad u \in \Phi(X), \quad i \in I \quad (2)$$

Let S consist of (1) and (2). Then S is a Gröbner-Shirshov basis in $RB(X)$.

Proof: Denote by $s \wedge l$ the composition of the polynomials of type s and type l .

The ambiguities of all possible compositions of the polynomials in S are only as below:

$$1 \wedge 1 \quad x_i x_j x_l, \quad i > j > l;$$

$$2 \wedge 1 \quad P(u)x_i x_j \text{ and } P(u)x_l = P(v|_{x_i x_j})x_l, \quad u \in \Phi(X), \quad v \in \Phi^*(X), \quad i > j;$$

$$2 \wedge 2 \quad P(u)x_i = P(v|_{P(w)x_j})x_i, \quad u, w \in \Phi(X), \quad v \in \Phi^*(X).$$

All possible compositions of left multiplication are $P(v) \cdot g$ where $v \in \Phi(X)$.

Now we check that all the compositions are trivial. We only check $1 \wedge 1$ as an example.

For $1 \wedge 1$, let $f = x_i x_j - x_j x_i, g = x_j x_l - x_l x_j$. Then $w = x_i x_j x_l$.

$$\begin{aligned} (f, g)_w &= (x_i x_j - x_j x_i)x_l - x_i(x_j x_l - x_l x_j) \\ &= x_i x_l x_j - x_j x_i x_l \\ &\equiv x_l x_j x_i - x_l x_j x_i \\ &\equiv 0 \quad \text{mod}(S, x_i x_j x_l). \end{aligned}$$

For any $P(v)$, let $P(v) = P^l(v')$ where $l \geq 1, v' \notin P(\Phi(X))$. Then

$$\begin{aligned} &P(v)(P(u)x_i - x_i P(u)) \\ &= P^l(v') \cdot (P^t(u')x_i - x_i P^t(u')) \\ &\equiv (P^l(v') \cdot P^t(u'))x_i - x_i(P^l(v') \cdot P^t(u')) \quad \text{mod}(S) \end{aligned}$$

where $l, t \geq 1, u', v' \notin P(\Phi(X))$. By Lemmas 3.1 and 3.2,

$$P^l(v') \cdot P^t(u') = \alpha_{(l,t,1)} P^{l+t-1}(P(v') \cdot u') + \varepsilon \triangleq \alpha_0 \varepsilon_0 + \sum_{j=1} \alpha_j \varepsilon_j$$

where $\varepsilon \in RB(X)$, each $\varepsilon_j \in P(\Phi(X))$, $\varepsilon_j < \overline{P^{l+t-1}(P(v') \cdot u')} = \overline{P^l(v') \cdot P^t(u')}$, $\alpha_j \in k, j = 1, 2, \dots$, $\varepsilon_0 = \overline{P^{l+t-1}(P(v') \cdot u')} = \overline{P^l(v') \cdot P^t(u')} \in P(\Phi(X))$ and $\alpha_0 = \alpha_{(l,t,1)}$. Therefore

$$P(v)(P(u)x_i - x_i P(u)) \equiv \alpha_0(\varepsilon_0 x_i - x_i \varepsilon_0) + \sum_{j=1} \alpha_j(\varepsilon_j x_i - x_i \varepsilon_j) \mod(S)$$

where each $\varepsilon_m x_i - x_i \varepsilon_m \in S$ and $\overline{\varepsilon_m x_i - x_i \varepsilon_m} \leq \overline{P^l(v') \cdot (P^t(u')x_i)} = \overline{P(v) \cdot (P(u)x_i)}, m \geq 0$. Thereby

$$P(v)(P(u)x_i - x_i P(u)) \equiv 0 \mod(S).$$

Thus all the compositions in S are trivial.

This shows that S is a Gröbner-Shirshov basis. ■

We now use the same notations as in Theorem 4.1. Clearly, $RB(X|S)$ is the free commutative Rota-Baxter algebra generated by X with weight λ .

The following corollary follows from our Theorems 3.9 and 4.1.

Corollary 4.2 ([38]) *Let $RB(X|S)$ be the free commutative Rota-Baxter algebra generated by X with weight λ as in Theorem 4.1. Then*

$$Irr(S) = Y_1 \cup Y_2 \cup Y_3$$

is a k -basis of $RB(X|S)$ where

$$\begin{aligned} Y_1 &= \{x_1 x_2 \cdots x_n \in S(X) | n \in \mathcal{N}^+, x_i \in X, x_1 \leq x_2 \leq \cdots \leq x_n\}, \\ Y_2 &= \{P^{l_1}(u_1 P^{l_2}(\cdots (u_{t-1} P^{l_t}(u_t)) \cdots)) \in \Phi(X) | u_j \in Y_1, t \geq 1, l_j \geq 1, j = 1, 2, \dots, t\}, \\ Y_3 &= \{uv \in \Phi(X) | u \in Y_1, v \in Y_2\}. \end{aligned}$$

■

Now, we show that every countably generated Rota-Baxter algebra with weight 0 can be embedded into a two-generated Rota-Baxter algebra.

Theorem 4.3 *Every countably generated Rota-Baxter algebra with weight 0 can be embedded into a two-generated Rota-Baxter algebra.*

Proof: Since every countably generated Rota-Baxter algebra A is countably dimensional, we may assume that A has a k -basis $X = \{x_i | i = 1, 2, \dots\}$. Then A can be expressed as $A = RB(X|S)$, where $S = \{x_i x_j = \{x_i, x_j\}, P(x_i) = \{P(x_i)\} \mid i, j \in \mathcal{N}^+\}$ and for any $u \in \Phi(X)$, $\{u\}$ means the linear combination of u by x_t , $x_t \in X$.

For $u \in \Phi(X)$, we denote $supp(u)$ the set of $x_i \in X$ appearing in the word u , for example, if $u = x_1 P(x_2) x_3 x_1$, then $supp(u) = \{x_1, x_2, x_3\}$.

Let $H = RB(X, a, b | S_1)$, where S_1 consists of the following relations:

- I. $x_i x_j = \{x_i x_j\}$, $i, j \in \mathcal{N}^+$,
- II. $P(x_i) = \{P(x_i)\}$, $i \in \mathcal{N}^+$,
- III. $aab^i ab = x_i$, $i \in \mathcal{N}^+$,
- IV. $P(t) = 0$, $t \in \Phi'(X, a, b)$,

where $\Phi'(X, a, b) = \{u \in \Phi(X, a, b) | \exists u' \in \Phi(X, a, b), u' \in Irr(\{III\}), supp(u') \cap \{a, b\} \neq \emptyset, s.t., u' \equiv u \mod(\{III\}, u)\}$.

Let the ordering on $\Phi(X, a, b)$ be defined as before, where $x_i > a > b$, $i \in \mathcal{N}^+$.

We will prove that S_1 is a Gröbner-Shirshov basis in $RB(X, a, b)$.

The ambiguities w of all possible compositions of in S_1 are:

$$1) \ x_i x_j x_l \quad 2) \ P(t|_{x_i x_j}) \quad 3) \ P(t|_{P(x_i)}) \quad 4) \ P(t|_{P(t')}) \quad 5) \ P(t|_{aab^i ab})$$

where $x_i, x_j, x_l \in X, t, t' \in \Phi'(X, a, b)$.

All possible compositions of left and right multiplications in S_1 are as below:

- 6) $P(u)(P(x_i) - \{P(x_i)\}), i \in \mathcal{N}^+, u \in \Phi(X, a, b)$,
- 7) $(P(x_i) - \{P(x_i)\})P(u), i \in \mathcal{N}^+, u \in \Phi(X, a, b)$,
- 8) $P(u)(P(t) - 0), u \in \Phi(X, a, b), t \in \Phi'(X, a, b)$,
- 9) $(P(t) - 0)P(u), u \in \Phi(X, a, b), t \in \Phi'(X, a, b)$.

We have to check that all these compositions are trivial. We just take 5) and 6) as examples.

For 5), let $f = P(t|_{aab^i ab})$, $g = aab^i ab - x_i$, $t \in \Phi'(X, a, b)$, $x_i \in X$. Then $w = P(t|_{aab^i ab})$ and

$$(f, g)_w = P(t|_{x_i}) \equiv 0 \mod(\{IV\}, w).$$

For 6),

$$\begin{aligned} & P(u)(P(x_i) - \{P(x_i)\}) \\ &= P(P(u)x_i) + P(uP(x_i)) - P(u)\{P(x_i)\} \\ &\equiv P(P(v)x_i) + P(vP(x_i)) - P(v)\{P(x_i)\} \mod(\{III\}) \end{aligned}$$

where $v \in \Phi(X, a, b)$ such that $u \equiv v \mod(\{III\}, u)$. There are two cases to consider.

a): If $v \in \Phi'(X, a, b)$, then $\text{mod}(\{IV\})$, $P(P(v)x_i) \equiv 0$, $P(vP(x_i)) \equiv 0$ and $P(v)\{P(x_i)\} \equiv 0$. Therefore,

$$P(u)(P(x_i) - \{P(x_i)\}) \equiv 0 \pmod{S_1}.$$

b): If $v \notin \Phi'(X, a, b)$, then $v \in \Phi(X)$. Since $v \equiv \sum \beta_j x_j \pmod{\{I, II\}}$,

$$P(u)(P(x_i) - \{P(x_i)\}) \equiv 0 \pmod{S_1}.$$

Thus, S_1 is a Gröbner-Shirshov basis in $RB(X, a, b)$. By Theroem 3.9, A can be embedded into the Rota-Baxter algebra H which is generated by $\{a, b\}$. ■

Recall that a dendriform dialgebra (see [45]) is a k -module D with two binary operations $\prec$ and $\succ$ such that

1. $(x \prec y) \prec z = x \prec (y \prec z + y \succ z)$
2. $(x \succ y) \prec z = x \succ (y \prec z)$
3. $(x \prec y + x \succ y) \succ z = x \succ (y \succ z)$

for any $x, y, z \in D$.

A dendriform trialgebra (see [46]) is a k -module T equipped with three binary operations $\prec, \succ$ and $\circ$ that satisfy the relations

1. $(x \prec y) \prec z = x \prec (y * z)$
2. $(x \succ y) \prec z = x \succ (y \prec z)$
3. $(x * y) \succ z = x \succ (y \succ z)$
4. $(x \succ y) \circ z = x \succ (y \circ z)$
5. $(x \prec y) \circ z = x \circ (y \succ z)$
6. $(x \circ y) \prec z = x \circ (y \prec z)$
7. $(x \circ y) \circ z = x \circ (y \circ z)$

for any $x, y, z \in T$ where $x * y = x \prec y + x \succ y + x \circ y$.

Denote by $D(X|S)$ ($T(X|S)$) the dendriform dialgebra (trialgebra) generated by X with defining relations S .

Suppose that $(D, \prec, \succ)$ is a dendriform dialgebra with a linear basis $X = \{x_n | n \in I\}$. Let $x_i \prec x_j = [x_i, x_j]$, $x_i \succ x_j = [[x_i, x_j]]$, where $[x_i, x_j]$ and $[[x_i, x_j]]$ are linear combinations of $x_n \in X$.

Then D has the expression

$$D = D(X|x_i \prec x_j = [x_i, x_j], x_i \succ x_j = [[x_i, x_j]]).$$

In [30], K. Ebrahimi-Fard and L. Guo proved that

$$U(D) = RB(X|x_i P(x_j) + \lambda x_i x_j = [x_i, x_j], P(x_i) x_j = [[x_i, x_j]])$$

is the universal enveloping algebra of D , where $\lambda \in k$.

Theorem 4.4 *Suppose that $(D, \prec, \succ)$ is a dendriform dialgebra with a linear basis $X = \{x_n | n \in I\}$, where I is a well ordered set. Let $x_i \prec x_j = [x_i, x_j]$, $x_i \succ x_j = [[x_i, x_j]]$, where $[x_i, x_j]$ and $[[x_i, x_j]]$ are linear combinations of $x_n \in X$. Let $\lambda \in k, \lambda \neq 0$. Let S be the set consisting of*

$$f_1 = x_i P(x_j) + \lambda x_i x_j - [x_i, x_j], \quad (3)$$

$$f_2 = P(x_i) x_j - [[x_i, x_j]], \quad (4)$$

$$f_3 = x_i x_j x_l - \lambda^{-1} [x_i, x_j] x_l + \lambda^{-1} x_i [[x_j, x_l]]. \quad (5)$$

Then in $RB(X)$, all intersection compositions and inclusion compositions in S are trivial.

Proof: The ambiguities of all possible compositions of the polynomials in S are only as below:

$$3 \wedge 4 \quad w_{34} = x_i P(x_j) x_l; \quad 4 \wedge 3 \quad w_{43} = P(x_i) x_j P(x_l);$$

$$4 \wedge 5 \quad w_{45} = P(x_l) x_i x_j x_s; \quad 5 \wedge 3 \quad w_{53} = x_i x_j x_s P(x_l);$$

$$5 \wedge 5 \quad w_{55} = x_i x_j x_s x_l \quad \text{and} \quad w'_{55} = x_i x_j x_h x_l x_s.$$

We only check $4 \wedge 3$ as example. The other cases are similar. Let $f = P(x_i) x_j - [[x_i, x_j]]$, $g = x_j P(x_l) + \lambda x_j x_l - [x_j, x_l]$. Then $w_{43} = P(x_i) x_j P(x_l)$ and

$$\begin{aligned} (f, g)_{w_{43}} &= (P(x_i) x_j - [[x_i, x_j]]) P(x_l) - P(x_i) (x_j P(x_l) + \lambda x_j x_l - [x_j, x_l]) \\ &= -\lambda P(x_i) x_j x_l + P(x_i) [x_j, x_l] - [[x_i, x_j]] P(x_l) \\ &\equiv -\lambda [[x_i, x_j]] x_l + [[x_i, [[x_j, x_l]]]] - (-\lambda [[x_i, x_j]] x_l + [[[x_i, x_j]], x_l]) \\ &\equiv -[[[x_i, x_j]], x_l] + [[x_i, [[x_j, x_l]]]] \\ &\equiv 0 \quad \text{mod}(S, P(x_i) x_j P(x_l)) \end{aligned}$$

by using the relation $(x \succ y) \prec z = x \succ (y \prec z)$.

Thus, the theorem is proved. ■

Suppose that $(T, \prec, \succ, \circ)$ is a dendriform trialgebra with a linear basis $X = \{x_n | n \in I\}$. Let $x_i \prec x_j = [x_i, x_j]$, $x_i \succ x_j = [[x_i, x_j]]$, $x_i \circ x_j = \langle x_i, x_j \rangle$, where $[x_i, x_j]$, $[[x_i, x_j]]$ and $\langle x_i, x_j \rangle$ are linear combinations of $x_n \in X$.

Then T has the expression

$$T = T(X | x_i \prec x_j = [x_i, x_j], x \succ x_j = [[x_i, x_j]], x_i \circ x_j = \langle x_i, x_j \rangle).$$

We notice in [30], K. Ebrahimi-Fard and L. Guo have proved that

$$U(T) = RB(X | x_i P(x_j) = [x_i, x_j], P(x_i) x_j = [[x_i, x_j]], \lambda x_i x_j = \langle x_i, x_j \rangle)$$

is the universal enveloping algebra of T , where $\lambda \in k$.

Theorem 4.5 *Suppose that $(T, \prec, \succ, \circ)$ is a dendriform trialgebra with a linear basis $X = \{x_n | n \in I\}$, where I is a well ordered set. Let $x_i \prec x_j = [x_i, x_j]$, $x_i \succ x_j = [[x_i, x_j]]$, $x_i \circ x_j = \langle x_i, x_j \rangle$, where $[x_i, x_j]$, $[[x_i, x_j]]$ and $\langle x_i, x_j \rangle$ are linear combinations of $x_n \in X$. Let S be the set consisting of*

$$f_4 = x_i P(x_j) - [x_i, x_j], \quad (6)$$

$$f_5 = P(x_i) x_j - [[x_i, x_j]], \quad (7)$$

$$f_6 = \lambda x_i x_j - \langle x_i, x_j \rangle. \quad (8)$$

Then in $RB(X)$, all intersection compositions and inclusion compositions in S are trivial.

Proof: The ambiguities of all possible compositions of the polynomials in S are only as below:

$$6 \wedge 7 \quad w_{67} = x_i P(x_j) x_l; \quad 7 \wedge 6 \quad w_{76} = P(x_i) x_j P(x_l);$$

$$7 \wedge 8 \quad w_{78} = P(x_l) x_i x_j; \quad 8 \wedge 6 \quad w_{86} = x_i x_j P(x_l);$$

$$8 \wedge 8 \quad w_{88} = \lambda x_i x_j x_l.$$

We only check $7 \wedge 6$ as example. The other cases are similar.

Let $f = P(x_i) x_j - [[x_i, x_j]]$, $g = x_j P(x_l) - [x_j, x_l]$. Then $w_{76} = P(x_i) x_j P(x_l)$ and

$$\begin{aligned} (f, g)_{w_{76}} &= (P(x_i) x_j - [[x_i, x_j]]) P(x_l) - P(x_i) (x_j P(x_l) - [x_j, x_l]) \\ &= P(x_i) [x_j, x_l] - [[x_i, x_j]] P(x_l) \\ &\equiv [[x_i, [x_j, x_l]]] - [[[x_i, x_j]], x_l] \\ &\equiv 0 \mod(S, P(x_i) x_j P(x_l)) \end{aligned}$$

by using the relation $(x \succ y) \prec z = x \succ (y \prec z)$.

Hence, the proof is completed. ■

References

- [1] William W. Adams and Philippe Loustau, An introduction to Gröbner bases, Graduate Studies in Mathematics, Vol. 3, American Mathematical Society (AMS), 1994.
- [2] Huihui An and Chengming Bai, From Rota-Baxter algebras to pre-Lie algebras. [arXiv:0711.1389v1]
- [3] M. Aquiar and M. Moreira, Combinatorics of the free Baxter algebra, *Electron J. Combin.*, **13**(1)(2006), to appear. [arXiv:math.CO/0510169]
- [4] F.V. Atkinson, Some aspects of Baxter's functional equation, *J. Math. Anal. Appl.*, **7**(1963), 1-30.
- [5] G. Baxter, An analytic problem whose solution follows from a simple algebraic identity, *Pacific J. Math.*, **10**(1960), 731-742.
- [6] Thomas Becker, Volker Weispfenning, and with cooperation from Heinz Kredel, Gröbner bases: A computational approach to commutative algebra, Graduate Texts in Mathematics, Vol.141, New York: Springer-Verlag, 1993.
- [7] G.M. Bergman, The diamond lemma for ring theory, *Adv. Math.*, **29**(1978), 178-218.
- [8] L.A. Bokut, Insolvability of the word problem for Lie algebras, and subalgebras of finitely presented Lie algebras, *Izvestija AN USSR (mathem.)*, **36**(6)(1972), 1173-1219.
- [9] L.A. Bokut, Imbeddings into simple associative algebras, *Algebra i Logika*, **15**(1976), 117-142.
- [10] L.A. Bokut and Yuqun Chen, Gröbner-Shirshov bases: Some new results, Proceedings of the Second International Congress in Algebra and Combinatorics, World Scientific, 2008, 35-56.
- [11] L.A. Bokut, Yuqun Chen and Cihua Liu, Gröbner-Shirshov bases for dialgebras. [arXiv:0804.0638]
- [12] L.A. Bokut, Yuqun Chen and Jianjun Qiu, Gröbner-Shirshov bases for associative algebras with multiple operators and free Rota-Baxter algebras, *Journal of Pure and Applied Algebra*, **214**(2010), 89-100.
- [13] L.A. Bokut, Y. Fong and W.-F. Ke, Composition Diamond lemma for associative conformal algebras, *J. Algebra*, **272**(2004), 739-774.

- [14] L.A. Bokut, Y. Fong, W.-F. Ke and P.S. Kolesnikov, Gröbner and Gröbner-Shirshov bases in algebra and conformal algebras, *Fundamental and Applied Mathematics*, **6**(3)(2000), 669-706.
- [15] L.A. Bokut and P.S. Kolesnikov, Gröbner-Shirshov bases: from their incipency to the present, *Journal of Mathematical Sciences*, **116**(1)(2003), 2894-2916.
- [16] L.A. Bokut and P.S. Kolesnikov, Gröbner-Shirshov bases, conformal algebras and pseudo-algebras, *Journal of Mathematical Sciences*, **131**(5)(2005), 5962–6003.
- [17] L.A. Bokut and G. Kukin, Algorithmic and Combinatorial algebra, Kluwer Academic Publ., Dordrecht, 1994.
- [18] B. Buchberger, An algorithmical criteria for the solvability of algebraic systems of equations [in German], *Aequationes Math.*, **4**(1970), 374-383.
- [19] B. Buchberger, G.E. Collins, R. Loos and R. Albrecht, Computer algebra, symbolic and algebraic computation, Computing Supplementum, Vol.4, New York: Springer-Verlag, 1982.
- [20] Bruno Buchberger and Franz Winkler, Gröbner bases and applications, London Mathematical Society Lecture Note Series, Vol.251, Cambridge: Cambridge University Press, 1998.
- [21] P. Cartier, On the structure of free Baxter algebras, *Adv. Math.*, **9**(1972), 253-265.
- [22] A. Connes and D. Kreimer, Hopf algebras, renormalization and noncommutative geometry, *Comm. in Math. Phys.*, **199**(1)(1998), 203-242.
- [23] A. Connes and D. Kreimer, Renormalization in quantum field theory and the Riemann-Hilbert problem. I. The Hopf algebra structure of graphs and the main theorem, *Comm. in Math. Phys.*, **210**(1)(2000), 249-273.
- [24] A. Connes and D. Kreimer, Renormalization in quantum field theory and the Riemann-Hilbert problem. II. The β -function, diffeomorphisms and the group, *Comm. in Math. Phys.*, **216**(1)(2001), 215-241.
- [25] David A. Cox, John Little and Donal O'Shea, Ideals, varieties and algorithms: An introduction to computational algebraic geometry and commutative algebra, Undergraduate Texts in Mathematics, New York: Springer-Verlag, 1992.
- [26] N.A. Derzko, Mappings satisfying Baxter's identity in the algebra of matrices, *J. Math. Anal. Appl.*, **42**(1973), 1-19.
- [27] K. Ebrahimi-Fard, Loday-type algebras and the Rota-Baxter relation, *Lett. Math. Phys.*, **61**(2)(2002), 139-147.

- [28] K. Ebrahimi-Fard, J. M. Gracia-Bondia and F. Patras, Rota–Baxter algebras and new combinatorial identities, *Lett. Math. Phys.*, **81**(2007), 61-65.
- [29] K. Ebrahimi-Fard and L. Guo, Rota-Baxter algebras in renormalization of perturbative quantum field theory, *Fields Inst. Commun.*, **50**(2007), 47-105.
- [30] K. Ebrahimi-Fard and L. Guo, Rota-Baxter algebras and dendriform algebras, *Journal of Pure and Applied Algebra*, **212**(2)(2008), 320-339.
- [31] K. Ebrahimi-Fard and L. Guo, Free Rota–Baxter algebras and rooted trees, *J. Algebra Applications*, **7**(2008), 167-194.
- [32] K. Ebrahimi-Fard and L. Guo, Multiple zeta values and Rota-Baxter algebras, *Integers*, Proceedings of CANT 2005 in Honor of Mel Nahtanson, **8**(2008).
- [33] K. Ebrahimi-Fard, L. Guo and D. Kerimer, Integrable renormalization II: the general case. [arXiv:hep-th/0403118v1]
- [34] David Eisenbud, Commutative algebra with a view toward algebraic geometry, Graduate Texts in Math., Vol.150, Berlin and New York: Springer-Verlag, 1995.
- [35] L. Guo, Baxter algebras and differential algebras, in “Differential algebra and related topics”, World Sci. Publishing Company, 2002.
- [36] L. Guo, Baxter Algebras, Stirling Numbers, and Partitions, *J. Algebra and Its Appl.*, **4**(2005), 153-164.
- [37] L. Guo and W. Keigher, Baxter algebras and Shuffle products, *Adv. Math.*, **150** (2000), 117-149.
- [38] L. Guo and W. Keigher, On free Baxter algebras: completions and the internal construction, *Adv. Math.*, **151** (2000), 101-127.
- [39] L. Guo and W. Keigher, On differential Rota-Baxter algebras, *Journal of Pure and Applied Algebra*, **212** (2008), 522-540.
- [40] L. Guo and William Y. Sit, Enumeration of Rota-Baxter words, Proceedings ISSAC 2006, Genova, Italy, ACM Press, 2006, 124-131. [arXiv:RA/0602449]
- [41] L. Guo and Bingyong Xie, Structure theorems of mixable shuffle algebras and free commutative Rota-Baxter algebras. [arXiv:0807.2267]
- [42] H. Hironaka, Resolution of singularities of an algebraic variety over a field of characteristic zero, I, II, *Ann. Math.*, **79**(1964), 109-203, 205-326.
- [43] S.-J. Kang and K.-H. Lee, Gröbner-Shirshov bases for irreducible sl_{n+1} -modules, *Journal of Algebra*, **232**(2000), 1-20.

- [44] X.X. Li, D.P. Hou and C.M. Bai, Rota-Baxter operators on pre-Lie algebras, *J. Nonlinear Math. Phys.*, **14**(2)(2007), 269-289.
- [45] J.-L. Loday, Dialgebras, in dialgebras and related operads, *Lecture Notes in Math.*, **1763**(2001), 7-66. (preprint 2001, arXiv:mathe.QA/0102053)
- [46] J.-L. Loday and M. Ronco, Algèbre de Hopf colibres, *C. R. Acad. Sci. Paris*, **337**(2003), 153-158.
- [47] S. Luiz, de Braganca, Finite dimensional Baxter algebras, *Studies in Applied Math.*, **54**(1)(1975), 75-89.
- [48] A.A. Mikhalev, The junction lemma and the equality problem for color Lie superalgebras, *Vestnik Moskov. Univ. Ser. I Mat. Mekh.*, **5**(1989), 88-91. English translation: *Moscow Univ. Math. Bull.*, **44**(1989), 87-90.
- [49] A.A. Mikhalev, The composition lemma for color Lie superalgebras and for Lie p -superalgebras, *Contemp. Math.*, **131**(2)(1992), 91-104.
- [50] A.A. Mikhalev, Shirshov's composition techniques in Lie superalgebra (non-commutative Gröbner bases). *Trudy. Sem. Petrovsk.*, **18**(1995), 277-289. English translation: *J. Math. Sci.*, **80**(1996), 2153-2160.
- [51] J.B. Miller, Some properties of Baxter operators, *Acta Math. Acad. Sci. Hungar.*, **17**(1966), 387-400.
- [52] J.B. Miller, Baxter operators and endomorphisms on Banach algebras, *J. Mat. Anal. Appl.*, **25**(1969), 503-520.
- [53] Nguyen Huu Bong, Some apparent connection between Baxter and averaging operators, *J. Math. Anal. Appl.*, **56**(2)(1976), 330-345.
- [54] G.-C. Rota, Baxter algebras and combinatorial identities I, II, *Bull. Amer. Math. Soc.*, **5**(1969), 325-329, 330-334.
- [55] G.-C. Rota, Baxter operators, an introduction, In: Gian-Carlo Rota on Combinatorics, Introductory papers and commentaries, J.P.S. Kung Ed., Contemp. Mathematicians, Birkhäuser Boston, Boston, MA, 1995.
- [56] G.-C. Rota, Ten mathematics problems I will never solve, *Mitt. Dtsch. Math-Ver.*, **2**(1998), 45-52.
- [57] A.I. Shirshov, Some algorithmic problem for ε -algebras, *Sibirsk. Mat. Z.*, **3**(1962), 132-137. (in Russian)
- [58] A.I. Shirshov, Some algorithmic problem for Lie algebras, *Sibirsk. Mat. Z.*, **3**(2)(1962), 292-296 (in Russian); English translation in SIGSAM Bull., **33**(2)(1999), 3-6.